

Before the Federal Communications Commission

Washington, D.C. 20554

In the Matter of

Reassessment of Federal Communications)	ET Docket No. 13-84
Commission Radiofrequency Exposure Limits and)	
Policies)	
)	
Proposed Changes in the Commission's Rules)	ET Docket No. 03-137
Regarding Human Exposure to Radiofrequency)	
Electromagnetic Fields)	

To: Office of the Secretary
Federal Communications Commission , Washington, DC 20554

As officially presented in the Federal Register/ Vol. 78, No. 107 / Tuesday, June 4, 2013 / Proposed Rules. Federal Communications Commission, 47 CFR Parts 1, 2, 15, 24, 25, 27, 73, 90, 95, 97, and 101 [ET Docket Nos. 03-137 and 13-84; FCC 13-39], Reassessment of Exposure to Radiofrequency Electromagnetic Fields Limits and Policies, Federal Communications Commission

Comment Filed by: Cindy Sage, Sage Associates
1396 Danielson Road, Santa Barbara, CA 93108
E-mail: sage@silcom.com Tel.: 805-969-0557

Prof. Lennart Hardell, MD
Department of Oncology Orebro University Hospital Orebro, Sweden
E-mail: lennart.hardell@orebroll.se

Dr. Martha Herbert, PhD, MD, Pediatric Neurology
TRANSCEND Research Program Massachusetts General Hospital
Harvard Medical School Boston, MA USA
Email: drmarthaherbert@gmail.com

1. New, biologically-based public exposure standards should be developed under the direction of experts in the biological effects and adverse health effects of chronic exposures to radiofrequency electromagnetic radiation (RFR), drawing upon the substantial international body of scientific and public health literature, and not be limited to individuals in electrical and electronic engineering.
2. A rapidly accumulating body of scientific evidence of harm to health and well-being constitute warnings that adverse health effects can occur with prolonged exposures to very low-intensity EMF at biologically active frequencies or frequency combinations.
3. The BioInitiative 2012 Report reports biological effects at exposure levels significantly below the 2007 recommended goal of 0.1 uW/cm². Since 2007, five new studies of base-station level RFR at intensities ranging from less than 0.001 uW/cm² to 0.05 uW/cm² report headaches, concentration difficulties and behavioral problems in children and adolescents; and sleep disturbances, headaches and concentration problems in adults. Exhibit A presents some representative studies (peer-reviewed and published in reputable scientific journals) that report biological effects and adverse health effects at levels that are clearly non-thermal (low-intensity). New biologically-based public exposure limits are critically needed in light of the vast rollout of wireless technologies that expose billions of people globally to elevated, artificial RFR (particularly pulsed RFR) in daily life. These studies are representative of several thousand studies over four decades that constitute emerging scientific evidence of risk to very low-intensity RFR with chronic exposure.
4. As new studies are completed and published on the effects of chronic, low-intensity RFR exposure across populations (from cell towers and wireless devices, for example) the results indicate adverse health impacts occur from on-going disruption of normal metabolism, endocrine function, male fertility parameters, fetal brain development, immune function, mental abilities, electrophysiology, and neural synchrony. Disruption of basic neural function due to artificial EMF/RFR exposures can disrupt weak-field effects that are necessary to guide non-linear biological oscillations and other cellular communications necessary for normal biological functioning, and result in unacceptable burdens on human health.

5. Evidence for Damage to Sperm and Reproduction

Evidence for damage to sperm and male reproduction parameters include adverse effects on sperm quality, motility and pathology in men who use and particularly those who wear a cell phone, PDA or pager on their belt or in a pocket (Agarwal et al, 2008; Agarwal et al, 2009; Wdowiak et al, 2007; De Iuliis et al, 2009; Fejes et al, 2005; Aitken et al, 2005; Kumar, 2012). Other studies conclude that usage of cell phones, exposure to cell phone radiation, or storage of a mobile phone close to the testes of human males affect sperm counts, motility, viability and structure (Aitken et al, 2004; Agarwal et al, 2007; Eroglu et al, 2006). Animal studies have demonstrated oxidative and DNA damage, pathological changes in the testes of animals, decreased sperm mobility and viability, and other measures of deleterious damage to the male germ line (Dasdag et al, 1999; Yan et al, 2007; Otitoloju et al, 2010; Salama et al, 2008; Behari et al, 2006; Kumar et al, 2012). There are fewer animal studies that have studied effects of cell phone radiation on female fertility parameters. Panagopoulous et al (2012) report decreased ovarian development and size of ovaries, and premature cell death of ovarian follicles and nurse cells in *Drosophila melanogaster*. Gul et al (2009) reported rats exposed to stand-by level RFR (phones on but not transmitting calls) had a decrease in the number of ovarian follicles in pups born to these exposed dams. Magras and Xenos (1997) reported irreversible infertility in mice after five (5) generations of exposure to RFR at cell phone tower exposure levels of less than one

microwatt per centimeter squared ($\mu\text{W}/\text{cm}^2$). See www.bioinitiative.org Section 18 for references.

HUMAN SPERM AND THEIR DNA ARE DAMAGED

Human sperm are damaged by cell phone radiation at very low intensities ($0.00034 - 0.07 \mu\text{W}/\text{cm}^2$). Many new studies in the last decade report sperm damage in humans and animals, leading to substantial concerns for fertility, reproduction and health of the offspring (unrepaired de novo mutations in sperm). Exposure levels are similar to those resulting from wearing a cell phone on the belt, or in the pants pocket, or using a wireless laptop computer on the lap. Sperm lack the ability to repair DNA damage.

6. Evidence for Brain Tumors

Based on epidemiological studies there is a consistent pattern of increased risk for glioma and acoustic neuroma associated with use of mobile phones and cordless phones. The evidence comes mainly from two study centres, the Hardell group in Sweden and the Interphone Study Group. No consistent pattern of an increased risk is seen for meningioma. A systematic bias in the studies that explains the results would also have been the case for meningioma. The different risk pattern for tumor type strengthens the findings regarding glioma and acoustic neuroma. Meta-analyses of the Hardell group and Interphone studies show an increased risk for glioma and acoustic neuroma. Supportive evidence comes also from anatomical localisation of the tumor to the most exposed area of the brain, cumulative exposure in hours and latency time that all add to the biological relevance of an increased risk. In addition risk calculations based on estimated absorbed dose give strength to the findings. See www.bioinitiative.org Section 11 for references.

- There is reasonable basis to conclude that RF-EMFs are bioactive and have a potential to cause health impacts.
- There is a consistent pattern of increased risk for glioma and acoustic neuroma associated with use of wireless phones (mobile phones and cordless phones) mainly based on results from case-control studies from the Hardell group and Interphone Final Study results.
- Epidemiological evidence gives that RF-EMF should be classified as a human carcinogen.
- The existing FCC/IEE and ICNIRP public safety limits and reference levels are not adequate to protect public health based on evidence for brain tumors and RFR exposure.
- New public health standards and limits are needed.

7. Evidence for Adverse Fetal and Neonatal Effects

Effects on the developing fetus from in-utero exposure to cell phone radiation have been observed in both human and animal studies since 2006. Sources of fetal and neonatal exposures of concern include cell phone radiation (both paternal use of wireless devices worn on the body and maternal use of wireless phones during pregnancy). Sources include exposure to whole-body RFR from base stations and WI-FI, use of wireless laptops, use of incubators for newborns with excessively high ELF-EMF levels resulting in altered heart rate variability and reduced melatonin levels in newborns, fetal exposures to MRI of the pregnant mother, and greater susceptibility to

leukemia and asthma in the child where there have been maternal exposures to ELF-EMF. Divan et al (2008) found that children born to mothers who used cell phones during pregnancy develop more behavioral problems by the time they have reached school age than children whose mothers did not use cell phones during pregnancy. Children whose mothers used cell phones during pregnancy had 25% more emotional problems, 35% more hyperactivity, 49% more conduct problems and 34% more peer problems (Divan et al, 2008). Aldad et al (2012) showed that cell phone radiation significantly altered fetal brain development and produced ADHD-like behavior in the offspring of pregnant mice. Exposed mice had a dose-dependent impaired glutamatergic synaptic transmission onto Layer V pyramidal neurons of the prefrontal cortex. The authors conclude the behavioral changes were the result of altered neuronal developmental programming in utero. Offspring mice were hyperactive and had impaired memory function and behavior problems, much like the human children in Divan et al (2008). Fragopoulou et al (2012) reports that brain astrocyte development followed by proteomic studies is adversely affected by DECT (cordless phone radiation) and mobile phone radiation. See www.bioinitiative.org Section 19 and 20 for references.

Fetal (in-utero) and early childhood exposures to cell phone radiation and wireless technologies in general may be a risk factor for hyperactivity, learning disorders and behavioral problems in school.

8. Evidence for Effects on Autism (Autism Spectrum Disorders)

*“Autism spectrum disorder (ASD), the fastest-growing complex neurodevelopment disorder, continues to rise in its prevalence, now affecting up to 1 in 50 children in the USA, and averaging 1% globally, according to the latest CDC report. More children will be diagnosed with ASD this year than with AIDS, diabetes & cancer combined in the USA. **ASD costs the nation \$137 billion a year and this debt is expected to increase in the next decade.** Hence, ASD has become a huge healthcare burden and global threat, categorized by the CDC as a national public health crisis.”* (Special Issue on Autism, North American Journal of Medicine and Science, Vol 6, Issue 3, July 2013, Harvard Medical School).

Several thousand scientific studies over four decades point to serious biological effects and health harm from EMF and RFR. These studies report genotoxicity, single-and double-strand DNA damage, chromatin condensation, loss of DNA repair capacity in human stem cells, reduction in free-radical scavengers (particularly melatonin), abnormal gene transcription, neurotoxicity, carcinogenicity, damage to sperm morphology and function, effects on behavior, and effects on brain development in the fetus of human mothers that use cell phones during pregnancy. Cell phone exposure has been linked to altered fetal brain development and ADHD-like behavior in the offspring of pregnant mice.

Many disrupted physiological processes and impaired behaviors in people with ASDs closely resemble those related to biological and health effects of EMF/RFR exposure. Biomarkers and indicators of disease and their clinical symptoms have striking similarities. At the cellular and molecular level many studies of people with ASDs have identified oxidative stress and evidence of free-radical damage, as well as deficiencies of antioxidants such as glutathione. Elevated intracellular calcium in ASDs can be associated with genetic mutations but more often may be downstream of inflammation or chemical exposures. Lipid peroxidation of cell membranes, disruption of calcium metabolism, altered brain wave activity and consequent sleep, behavior and

immune disfunction, pathological leakage of critical barriers between gut and blood or blood and brain may also occur. Mitochondria may function poorly, and immune system disturbances of various kinds are common. Changes in brain and autonomic nervous system electrophysiology can be measured and seizures are far more common in ASCs than in the population at large. Sleep disruption and high levels of stress are close to universal in ASCs. All of these phenomena have also been documented to result from or be modulated by EMF/RFR exposure. Reducing or removing EMF and wireless RFR stressors from the environment is a reasonable precautionary action given the overall weight of evidence for a link to ASDs. The FCCs thermal safety limits do not address low-intensity (non-thermal) effects. The evidence is now overwhelming that limiting exposures to those causing thermal injury alone does not address the much broader array of risks and harm now clearly evident with chronic exposure to low-intensity (non-thermal) EMF/RFR. The now well-documented genotoxic impacts of EMF/RFR, placed in parallel with the huge rise in reported cases of ASCs as well as with the de novo mutations associated with some cases of ASCs (as well as other conditions), make it urgent to address the issue of (environmental) acquired as well as inherited genetic damage. With the rising numbers people with ASCs and other childhood health and developmental disorders, and with emerging evidence that EMF/RFR is a preventable environmental exposure of consequence to ASCs; public safety limits must be rethought in terms of fetal, neonatal and childhood neurological and electrophysiological development. The evidence is sufficient to warrant new public exposure standards benchmarked to low-intensity (non-thermal) exposure levels causing biological disruption and strong, interim precautionary practices are advocated. See www.bioinitiative.org Section 20 for references.

9. FCC Dockets 13-84, 03-137 and 13-39 propose to significantly relax rather than tighten exposure standards, in stark contrast to what the scientific evidence suggests is needed to protect public health from RFR. IEEE/FCC public safety limits remain unchanged and are still inadequate and obsolete with respect to prolonged, low-intensity NIER exposures.

Respectfully submitted:

Cindy Sage, Sage Associates 1396 Danielson Road, Santa Barbara, CA 93108
E-mail: sage@silcom.com Tel.: 805-969-0557

Prof. Lennart Hardell, MD Department of Oncology
Orebro University Hospital Orebro, Sweden
E-mail: lennart.hardell@orebroll.se

Dr. Martha Herbert, PhD, MD, Pediatric Neurology
TRANSCEND Research Program
Massachusetts General Hospital
Harvard Medical School Boston, MA USA
Email: drmarthaherbert@gmail.com

Exhibit A

Reported Biological Effects from Radiofrequency Radiation at Low-Intensity Exposure (Cell Tower, Wi-Fi, Wireless Laptop and 'Smart' Meter RF Intensities (Pages 1 – 11)

<http://www.bioinitiative.org/rf-color-charts/>

 [DOWNLOAD RF Color Charts](#)



[DOWNLOAD RF Color Charts](#)

Reported Biological Effects from Radiofrequency Radiation at Low-Intensity Exposure (Cell Tower, Wi-Fi, Wireless Laptop and 'Smart' Meter RF Intensities)

Power Density (Microwatts/centimeter ² - uW/cm ²)	Reference
As low as (10 ⁻¹³) or 100 femtowatts/cm ²	Belyaev, 1997
5 picowatts/cm ² (10 ⁻¹²)	Grundler, 1992
0.1 nanowatt/cm ² (10 ⁻¹⁰) or 100 picowatts/cm ²	Belyaev, 1997
0.00034 uW/cm ²	Behari, 2006
0.0005 uW/cm ²	Velizarov, 1999
0.0006 - 0.001 uW/cm ²	Buchner, 2012
0.0006 - 0.0128 uW/cm ²	Oberfeld, 2004
0.0009 uW/cm ²	Stagg, 1997
0.003 - 0.02 uW/cm ²	Heinrich, 2010
0.003 to 0.05 uW/cm ²	Thomas, 2010
0.005 uW/cm ²	Mohler, 2010
0.005 - 0.04 uW/cm ²	Thomas, 2008
0.01 - 0.11 uW/cm ²	Navarro, 2003

Stress proteins, HSP, disrupted immune function	Brain tumors and blood-brain barrier
Reproduction/fertility effects	Sleep, neuron firing rate, EEG, memory, learning, behavior
Oxidative damage/ROS/DNA damage/DNA repair failure	Cancer (other than brain), cell proliferation
Disrupted calcium metabolism	Cardiac, heart muscle, blood-pressure, vascular effects

Reference List
Reported Biological Effects from Radiofrequency Radiation (RFR)
at Low-Intensity Exposure Levels

(Cell Tower, WI-FI, Wireless Laptop, Wireless Utility Meters 'smart meters')

<http://www.bioinitiative.org/bibliography/>  [DOWNLOAD REFERENCE LIST \(PDF\)](#)

Acherman P et al, 2000. Exposure to pulsed high-frequency electromagnetic field during waking affects human sleep EEG. *NeuroReport* 11(15):3321-3325.

Adey,WR et al, 1999. Incidence of spontaneous and nitrosourea-induced primary tumors of the central nervous system in Fischer 344 rats chronically exposed to modulated microwaves. *Radiation Research* 152: 293-302.

Agarwal A, Deepinder F, Sharma RK, Ranga G, Li J. 2008. Effect of cell phone usage on semen analysis in men attending infertility clinic: an observational study. *Fertil Steril.* 89(1):124-8.

Agarwal A, Desai NR, Makker K, Varghese A, Mouradi R, Sabanegh E, Sharma R. 2009. Effects of radiofrequency electromagnetic waves (RF-EMW) from cellular phones on human ejaculated semen: an in vitro pilot study. *Fertil Steril.* 92(4) 1318-1325.

Aitken RJ, Bennetts LE, Sawyer D, Wiklendt AM, King BV. 2005 Impact of radio frequency electromagnetic radiation on DNA integrity in the male germline 28:171-179.

Akoev, IG et al, 2002. Enzymatic activity of some tissues and blood serum from animals and humans exposed to microwaves and hypothesis on the possible role of free radical processes in the nonlinear effects and modification of emotional behavior of animals. *Radiats Biol Radioecol*, 42(3):32-330.

Atasoy HI, Gunal MY, Atasoy P, Elgun S, Bugdayci G. 2012 Immunohistopathologic demonstration of deleterious effects on growing rat testes of radiofrequency waves emitted from conventional Wi-Fi devices. *J Pediatr Urol.* [Epub ahead of print]

Avendano C, Mata A, Sanchez Sarmiento CA, Doncei GF. 2012. Use of laptop computers connected to internet through Wi-Fi decreases human sperm motility and increases sperm DNA fragmentation. *Fertility and Sterility.* American Society for Reproductive Medicine, Published by Elsevier Inc. doi:10.1016/j.fertnstert.2011.10.012.

Beason, RC & Semm, P, 2002. Responses of neurons to an amplitude modulated microwave stimulus. *Neuroscience Letters* 333:175-178.

Behari J, Kesari KK 2006. Effects of microwave radiations on reproductive system of male rats. *Embryo Talk* 1 (Suppl.1):81-5.

Belokrinsky, VS, 1982,. Destructive and reparative processes in hippocampus with long-term exposure to nonionizing radiation. In: U.S.S.R. Report, Effects of Nonionizing Microwave Radiation, No. 7, JPRS 81865, pp. 15-20.

- Belyaev IY, Alipov YD, Harms-Ringdahl M. 1997. Effects of zero magnetic field on the conformation of chromatin in human cells. *Biochim Biophys Acta* 1336(3):465-473.
- Belyaev IY, Hillert L, Protopopova M, Tamm C, Malmgren LO, Persson BR, Selivanova G, Harms-Ringdahl M. 2005. 915 MHz microwaves and 50 Hz magnetic field affect chromatin conformation and 53BP1 foci in human lymphocytes from hypersensitive and healthy persons. *Bioelectromagnetics*. 26(3):173-184.
- Belyaev IY, Markova E, Hillert L, Malmgren LOG, Persson BRR. 2009. Microwaves from UMTS/GSM mobile phones induce long-lasting inhibition of 53BP1/ γ -H2AX DNA repair foci in human lymphocytes. *Bioelectromagnetics* 30(2):129-41.
- Bolshakov, MA & Alekseev, SI, 1992. Bursting responses of Lymnea neurons to microwave radiation. *Bioelectromagnetics* 13(2): 119-129.
- Borbely, AA et al, 1999. Pulsed high-frequency electromagnetic field affects human sleep and sleep electroencephalogram. *Neuroscience Letters* 275(3): 207-210.
- Boscolo et al, 2001. Effects of electromagnetic fields produced by radiotelevision broadcasting stations on the immune system of women. *Sci Total Environ* 273(1-3):1-10.
- Buchner K, Eger H., 2011. Changes of Clinically Important Neurotransmitters under the Influence of Modulated RF Fields—A Long-term Study under Real-life Conditions *Umwelt-Medizin-Gesellschaft* 24(1): 44-57. Original study in German.
- Chia SE et al, 2000. Prevalence of headache among handheld cellular telephone users in Singapore: A Community Study. *Environmental Health Perspectives* 108(11):1059-1062.
- Chiang, H et al, 1989. Health effects of environmental electromagnetic fields. *Journal of Bioelectricity* 8: 127-131.
- Chou, CK et al, 1992. Long-term low level microwave irradiation of rats. *Bioelectromagnetics* 13:469-496.
- Czyz J et al, 2004. High frequency electromagnetic fields (GSM signals) affect gene expression levels in tumor suppressor p53-deficient embryonic stem cells. *Bioelectromagnetics* 25: 296-307.
- Dasdag, S et al, 1999. Whole-body microwave exposure emitted by cellular phones and testicular function of rats. *Urological Research* 27(3):219-223.
- D'Costa, H et al. 2003. Human brain wave activity during exposure to radiofrequency field emissions from mobile phones. *Australasian Physical & Engineering Sciences in Medicine*, Vol. 26, No. 4
- De Pomerai, D et al, 2000. Non-thermal heat-shock response to microwaves. *Nature* 405: 417-418.
- D'Inzeo, G et al, 1988. Microwave effects on acetylcholine-induced channels in cultured chick myotubes. *Bioelectromagnetics* 9: 363-372.
- De Iuliis GN, Newey RJ, King BV, Aitken RJ. 2009. Mobile phone radiation induces reactive oxygen species production and DNA damage in human spermatozoa in vitro. *PLoS One* 4(7):e6446.

Dolk, H et al, 1997. Cancer incidence near radio and television transmitters in Great Britain. *American Journal of Epidemiology* 145(1): 1-9.

Dumansky, JD & Shandala, MG, 1974. The biological action and hygienic significance of electromagnetic fields of superhigh and ultrahigh frequencies in densely populated areas. In: *Biological Effects and Health Hazards of Microwave Radiation. Proceedings of an International Symposium*, Czernski, P et al, (Eds) Warsaw, 15-18 October 1973, Polish Medical Publishers.

Dutta, SK et al, 1989. Radiofrequency radiation-induced calcium ion efflux enhancement from human and other neuroblastoma cells in culture. *Bioelectromagnetics* 10: 197-202.

Elekes, E, 1996. Effect on the immune system of mice exposed chronically to 50 Hz amplitude-modulated 2.45 GHz microwaves. *Bioelectromagnetics* 17:246-248.

Eltiti S, Wallace D, Ridgewell A, Zougkou K, Russo R, Sepulveda F, Mirshekar-Syahkal D, Rasor P, Deeble R, Fox E. 2007. Does short-term exposure to mobile phone base station signals increase symptoms in individuals who report sensitivity to electromagnetic fields? A double-blind randomized provocation study. *Environ Health Perspect* 115(11): 1603-8.

Eltiti S, Wallace D, Ridgewell A, Zougkou K, Russo R, Sepulveda F, Fox E. 2009. Short-term exposure to mobile phone base station signals does not affect cognitive functioning or physiological measures in individuals who report sensitivity to electromagnetic fields and controls. *Bioelectromagnetics* 30(7):556-63.

Federal Communications Commission, 1997. OET Bulletin 65: 1997-01

Fesenko, EE et al, 1999. Microwaves and cellular immunity. I. Effect of whole body microwave irradiation on tumor necrosis factor production in mouse cells. *Bioelectrochemistry and Bioenergetics* 49 (1): 29-35.

Fragopoulou AF, Koussoulakos SL, Margaritis LH. 2010. Cranial and postcranial skeletal variations induced in mouse embryos by mobile phone radiation. *Pathophysiology*. 17(3):169-77.

Garaj-Vrhovac, V et al, 1999. Micronucleus assay and lymphocyte mitotic activity in risk assessment of occupational exposure to microwave radiation. *Chemosphere* 39 (13) 2301-2312.

Grundler W, Kaiser F, Keilmann F, Walleczek J. 1992. Mechanisms of electromagnetic interaction with cellular systems. *Naturwissenschaften* 79(12):551-9.

Gul A, Celebi H, Uğraş S. 2009. The effects of microwave emitted by cellular phones on ovarian follicles in rats. *Arch Gynecol Obstet*. 280(5):729-33,

Hamblin, D. et al, 2004. Examining the effects of electromagnetic fields emitted by GSM mobile phones on human event-related potentials and performance during an auditory task. *Clinical Neurophysiology* 115:171-178.

Hamnerius, Y, 2000. Microwave exposure from mobile phones and base stations in Sweden. International Conference on Cell Tower Siting, June 7-8, 2000, Sponsored by the University of Vienna & Land Salzburg, Salzburg, Austria.

- Heinrich S, Thomas S, Heumann C, von Kries R, Radon K. 2010. Association between exposure to radiofrequency electromagnetic fields assessed by dosimetry and acute symptoms in children and adolescents: a population based cross-sectional study. *Environ Health* 9:75.
- Hjollund NH, Bonde JP, Skotte J, 1997 Semen analysis of personnel operating military radar equipment. *Reprod Toxicol* 11(6):897
- Hocking, B et al, 1996. Cancer incidence and mortality and proximity to TV towers *Medical Journal of Australia* 165(11-12): 601-605.
- Hocking, B et al, 2000. Decreased survival for childhood leukemia in proximity to TV towers. Poster presented at the Annual Scientific Meeting of the Royal Australian College of Physicians in Adelaide, SA, Australia, May 2000.
- Huber, R et al, 2002. Electromagnetic fields, such as those from mobile phones alter regional cerebral blood flow and sleep and waking EEG. *J. Sleep Res.* 11: 289-295.
- Hutter HP, Moshhammer H, Wallner P, Kundi M. 2006. Subjective symptoms, sleeping problems, and cognitive performance in subjects living near mobile phone base stations, *Occup. Environ. Med.* 63. 307–313.
- IEEE, 1999. C95: 1-1999(US)
- Ilhan A, Gurel A, Armutcu F, Kamisli S, Iraz M, Akyol O, Ozen S. 2004, Ginkgo biloba prevents mobile phone-induced oxidative stress in rat brain. *Clin Chim Acta.* 340(1-2): 153-162.
- Ilhan, A et al. 2004. Ginkgo biloba prevents mobile phone-induced oxidative stress in rat brain. *Clinica Chimica Acta* 340, 153-162.
- Ivaschuk, OI et al, 1997. Exposure of nerve growth factor-treated PC 12 rat pheochromocytoma cells to a modulated radiofrequency field at 836.55 MHz: effects on c-jun and c-fos expression. *Bioelectromagnetics* 18 (3): 223-229.
- Kesari KK, Behari J. 2012 Evidence for mobile phone radiation exposure effects on reproductive pattern of male rats: Role of ROS. *Electromagn Biol Med.* 31(3):213-22.
- Kong X, McDougale CJ (Editors). Preface. Special Issue on Autism, *North American Journal of Science and Medicine*, July 2013 (Volume 6, Issue 3)
<http://najms.net/wp-content/uploads/v06i03.pdf>
- Khurana VG, Hardell L, Everaert J, Bortkiewicz A, Carlberg M, Ahonen M. 2010. Epidemiological evidence for a health risk from mobile phone base stations. *Int J Occup Environ Health.* 16(3):263-267.
- Koivisto, M et al, 2000a. Effects of 902 MHz electromagnetic field emitted by cellular telephones on response times in humans. *Neuroreport* 11: 413-415.
- Koivisto, M et al, 2000b. The effects of electromagnetic field emitted by GSM phones on working memory. *Neuroreport* 11:1641-1643.

Kolodynski, AA, & Kolodynska VV, 1996. Motor and psychological functions of school children living in the area of the Skrunda radio location station in Latvia. *Science of the Total Environment* 180:87-93.

Krause, CM et al, 2000. Effects of electromagnetic field emitted by a cellular phone on the EEG during a memory task. *Neuroreport* 11:761-764.

Kues, HA et al, 1992. Increased sensitivity of the non-human primate eye to radiation following ophthalmic drug pretreatment. *Bioelectromagnetics* 13:379-393.

Kumar S Behari J Sisodia R. 2012. Impact of Microwave at X-Band in the aetiology of male infertility. *Electromagnetic Biology and Medicine*, 31(3): 223–232. online DOI: 10.3109/15368378.2012.700293.

Kundi M. Hutter HP. 2009. Mobile phone base stations—Effects on wellbeing and health. *Pathophysiology* 16 123–135.

Kwee, S et al, 1997. The biological effects of microwave radiation. *Proceedings of the Second World Congress for Electricity and Magnetism in Biology and Medicine*, Bologna, Italy, June 1997.

Kwee, S et al, 2001. Changes in cellular proteins due to environmental non-ionizing radiation. I. Heat-shock proteins. *Electro-and Magnetobiology* 20:141-152.

Lai H, & Singh, NP, 1996. Single and double strand DNA breaks in rat brain cells after acute exposure to radiofrequency electromagnetic radiation. *International Journal of Radiation Biology* 69:513-21.

Lass, J et al, 2002. Effects of 7 Hz-modulated 450 MHz electromagnetic radiation on human performance in visual memory tasks. *Int. J. Radiat. Biol.* 73(10): 937-944.

Lebedeva NN et al, 2000. Cellular phone electromagnetic field effects on bioelectric activity of human brain. *Crit Rev Biomed Eng* 28(1-2) 323-337.

Leszczynski, D et al, 2002. Non-thermal activation of the hsp27/p38MAPK stress pathway by mobile phone radiation in the human endothelial cells: Molecular mechanism for cancer- and blood-brain barrier-related effects. *Differentiation* 70: 120-129.

Leszczynski, D et al, 2004. Proteomics analysis of human endothelial cell line EA.hy926 after exposure to GSM 900 radiation. *Short Communication. Proteomics* 4, 1359-1365.

Lu, ST et al, 1999. Ultrawide-band electromagnetic pulses induced hypotension in rats. *Physiology and Behavior*, 67:753-761.

Magras, IN & Zenos, TD, 1997. RF Radiation-induced changes in the prenatal development of mice. *Bioelectromagnetics* 18:455-461.

Mann, K et al, 1996. Effects of pulsed high-frequency electromagnetic fields on human sleep. *Neuropsychobiology* 33:41-47.

Mann, K et al, 1998. Effects of pulsed high-frequency electromagnetic fields on the neuroendocrine system. *Neuroendocrinology* 67: 139-144.

Mantiply, ED et al, 1997. Summary of measured radiofrequency electric and magnetic fields (10 kHz to 30 GHz) in the general and work environment. *Bioelectromagnetics* 18: 563-577.

Markova E, Hillert L, Malmgren L, Persson BRR, Belyaev IY. 2005. Microwaves from GSM mobile telephones affect 53BP1 and γ -H2AX foci in human lymphocytes from hypersensitive and healthy persons. *Environmental Health Perspectives* Vol 113: No. 91 1172-1177

Marinelli, F La Sala D Cicciotti G Cattini L Trimarchi C Putti S Zamparelli A Giuliani L Tomassetti G Cinti C. 2004. Exposure to 900 MHz Electromagnetic Field induces an unbalance between pro-apoptotic and pro-survival signals in T-lymphoblastoid leukemia CCRF-CEM cells. *Journal of Cellular Physiology* 198: 324 – 332.

Mohler E, Frei P, Braun-Fahrlander C, Fröhlich J, Neubauer G, Rösli M; Qualifex Team. 2010. Effects of everyday radiofrequency electromagnetic-field exposure on sleep quality: a cross-sectional study. *Radiat Res* 174(3):347-56.

Oberfeld, G Enrique, NA Manuel P Ceferino M Gomez-Perretta C. 2004. The Microwave Syndrome – Further Aspects of a Spanish Study. 3rd International Workshop on Biological Effects of Electromagnetic Fields. Kos, Greece.

Otitolaju AA, Obe IA, Adewale OA, Otubanjo OA, Osunkalu VO. 2010. Preliminary study on the induction of sperm head abnormalities in mice, *Mus musculus*, exposed to radiofrequency radiations from global system for mobile communication base stations. *Bulletin of Environmental Contamination and Toxicology* 84(1):51-4.

Navakatikian, MA & Tomashevskaya, LA, 1994 Phasic behavioral and endocrine effects of microwaves of nonthermal intensity. In: *Biological Effects of Electric and Magnetic Fields*, Volume 1, Carpenter, DO, (Ed.) Academic Press, Inc., San Diego, CA., pp. 333-342.

Navarro EA, Sequra J, Portoles M, Gomez-Perretta de Mateo C. 2003. The Microwave Syndrome: A Preliminary Study in Spain. *Electromag Biol Med* 22:161-169,

Novoselova, EG et al, 1999. Microwaves and cellular immunity. II Immunostimulating effects of microwaves and naturally occurring antioxidant nutrients. *Bioelectrochemistry and Bioenergetics* 49 (1): 37-41.

Panagopoulos DJ. 2012. Effect of microwave exposure on the ovarian development of *Drosophila melanogaster*. *Cell Biochem Biophys*. 63(2):121-132,.

Paulraj R, Behari J. 2004. Radio frequency radiation effects on protein kinase C activity in rats' brain. *Mutat Res*. 545(1-2):127-130,

Persson, RR et al, 1997. Blood-brain barrier permeability in rats exposed to electromagnetic fields used in wireless communication. *Wireless Networks* 3:455-461.

Paredi P et al, 2001. Local Vasodilator Response to Mobile Phones. *Laryngoscope* 111: 159-162.

Phillips, J et al, 1998. DNA damage in molt-4 lymphoblastoid cells exposed to cellular telephone radiofrequency fields in vitro. *Bioelectrochemistry and Bioenergetics* 45:103-110.

Preece, A et al, 1999. Effect of a 915-MHz simulated mobile phone signal on cognitive function in man. *International Journal of Radiation Biology* 75: 447-456.

Pyrpasopoulou, A et al, 2004. Bone morphogenetic protein expression in newborn rat kidneys after prenatal exposure to radiofrequency radiation. *Bioelectromagnetics* 25: 216-227.

Ray, S & Behari, J, 1990. Physiological changes in rats after exposure to low levels of microwaves. *Radiation Research* 123: 190-202.

Repacholi, M. et al, 1997. Lymphomas in *Eμ-Pim1* transgenic mice exposed to pulsed 900 MHz electromagnetic fields. *Radiation Research* 147:31-40.

Richter, E et al, 2000. Cancer in radar technicians exposed to radiofrequency/microwave radiation: sentinel episodes. *International Journal of Occupational Health* 6(3): 187-193.

Riddervold IS, Pedersen GF, Andersen NT, Pedersen AD, Andersen JB, Zachariae R, Mølhave L, Sigsgaard T, Kjaergaard SK. 2008. Cognitive function and symptoms in adults and adolescents in relation to RF radiation from UMTS base stations. *Bioelectromagnetics* 29(4):257-67.

Sage Associates, 2004. An Overview of Low-Intensity Radiofrequency/Microwave Radiation Studies Relevant to Wireless Communications and Data. *Bioelectromagnetics Society Annual Meeting*, Washington DC, June 2004.

Sage Associates, 2004. Epidemiology for Decisionmakers: A Visual Guide to Residential and Occupational EMF Epidemiological Results on Leukemia 1979-2004. International Conference on Leukemia, London, September 2004. Children with Leukemia Trust (UK Registered Charity No. 298405).

Sage Associates, 2000. An overview of radiofrequency/microwave radiation studies relevant to wireless communications and data. International Conference on Cell Tower Siting, Salzburg, Austria, Land Salzburg-Landessanitätsdirektion – Umweltmedizin, Federal State of Salzburg Public Health Department, Environmental Health Unit, June 7-8, 2000.

Salama N, Kishimoto T, Kanayama HO. 2010. Effects of exposure to a mobile phone on testicular function and structure in adult rabbit. *Int J Androl.* 33(1):88-94.

Salford, LG et al. 1994. Permeability of the blood brain barrier induced by 915 MHz electromagnetic radiation, continuous wave and modulated at 8, 16, 50 and 200 Hz. *Microscopy Research and Technique* 27:535-542.

Salford, LG et al, 2003. Nerve cell damage in mammalian brain after exposure to microwaves from GSM mobile phones. *Environmental Health Perspectives Online* January 29.

Sandstrom M et al, 2001. Mobile phone use and subjective symptoms. Comparison of symptoms experienced by users of analogue and digital mobile phones. *Occup Med* 51(1) 25-35.

Santini R et al, 2001. Symptoms rapportes par des utilisateurs de telephones mobiles cellulaires. *Path Biol* 49:222-226.

Sarimov, R., Malmgren, L.O.G., Markova, E., Persson, B.R.R., Belyaev, I.Y. 2004. Nonthermal GSM microwaves affect chromatin conformation in human lymphocytes similar to heat shock. *IEEE Trans Plasma Sci* 32:1600-1608,

Schirmacher, A et al, 2000. Electromagnetic fields (1.8 GHz) increase the permeability of sucrose of the blood-brain barrier in vitro. *Bioelectromagnetics* 21:338-345.

Schmidt M Murbach M Inelustenberger C Maire M Kuster N Achermann P. s2012. Sleep EEG alterations: effects of pulsed magnetic fields versus pulse-modulated radio frequency electromagnetic fields. *J. Sleep Research, European Sleep Research Society* DOI: 10.1111/j.1365-2869.2012.01025.x.

Schwartz, JL et al, 1990. Exposure of frog hearts to CW or amplitude-modulated VHF fields: selective efflux of calcium ions at 16Hz. *Bioelectromagnetics* 11(4): 349-358.

Seaman, RL et al, 1999. Hyperactivity caused by nitric oxide synthase inhibitor is countered by ultra-wide band pulses. *Bioelectromagnetics* 20: 431-439.

Somogyi, Z et al, 1991. Effects of modulated and continuous microwave irradiation on the morphology and cell surface negative charge of 3T3 fibroblasts. *Scanning Microsc* 5(4):1145-1155.

Somogyi, Z et al, 1993. Effects of modulated and continuous microwave irradiation on pyroantimonate precipitable calcium content in junctional complex of mouse small intestine. *Scanning Microsc* 7(4): 1255-1261.

Stagg RB et al, 1997. DNA synthesis and cell proliferation in C6 glioma and primary glial cells exposed to 836.55 MHz modulated radiofrequency field. *Bioelectromagnetics* 18(3):230-236.

Stang A et al, 2001. The possible role of radiofrequency radiation in the development of uveal melanoma. *Epidemiology* 12(1):7-12.

Stankiewicz W, Dąbrowski MP, Kubacki R, Sobiczewska E, Szmigielski S. 2006. Immunotropic Influence of 900 MHz Microwave GSM Signal on Human Blood Immune Cells Activated in Vitro. *Electromagnetic Biology and Medicine* 25(1) 45-51.

Stark KD et al, 1997. Absence of chronic effect of exposure to short-wave radio broadcast signal on salivary melatonin concentrations in dairy cattle. *J Pineal Res* 22(4):171:176.

Sun W, Shen X, Lu D, Fu Y, Chiang H. 2012. A 1.8-GHz radiofrequency radiation induces EGF receptor clustering and phosphorylation in cultured human amniotic (FL) cells. *Int J Radiat Biol* 88(3):239-44.

Szmigielski, S et al, 1982. Accelerated development of spontaneous and benzpyrene-induced skin cancer in mice exposed to 2350 MHz microwave radiation. *Bioelectromagnetics* 3: 179-192.

Tattersall, JE et al, 2001. Effects of low intensity radiofrequency electromagnetic fields on electrical activity in rat hippocampal slices. *Brain Res* 904(1): 43-53.

Thomas S, Kühnlein A, Heinrich S, Praml G, Nowak D, von Kries R, Radon K. 2008. Personal exposure to mobile phone frequencies and well-being in adults: a cross-sectional

study based on dosimetry. *Bioelectromagnetics* 29:463-470.

Thomas S, Heinrich S, von Kries R, Radon K. 2010. Exposure to radio-frequency electromagnetic fields and behavioural problems in Bavarian children and adolescents. *Eur J Epidemiol* 25(2):135-41.

TNO Physics and Electronics Laboratory, The Netherlands. 2003. Effects of Global Communication System radio-frequency fields on well-being and cognitive functions of human beings with and without subjective complaints. Netherlands Organization for Applied Scientific Research 1-63.

Trosic, I et al, 2002. Micronucleus induction after whole-body microwave irradiation of rats. *Mutation Research* 521: 73-79.

Velizarov, S et al, 1999. The effects of radiofrequency fields on cell proliferation are non-thermal. *Bioelectrochemistry and Bioenergetics* 48: 177-180.

Veyret, B et al, 1991. Antibody responses of mice exposed to low-power microwaves under combined, pulse and amplitude modulation. *Bioelectromagnetics* 12: 47-56.

Weisbrot, D et al, 2003. Effects of mobile phone radiation on reproduction and development in *Drosophila melanogaster*. *Journal of Cellular Biochemistry* 89: 48-55.

Wolke, S et al, 1996. Calcium homeostasis of isolated heart muscle cells exposed to pulsed high-frequency electromagnetic fields. *Bioelectromagnetics* 17(2): 144-153.

Yan JG, Agresti M, Bruce T, Yan YH, Granlund A, Matloub HS. 2007. Effects of cellular phone emissions on sperm motility in rats. *Fertility and Sterility* 88(4):957-64.